

Non-Gravitating Scalar Field in the FRW Background

Nabamita Banerjee,^{*} Rajeev Kumar Jain,[†] and Dileep P. Jatkar[‡]
Harish-Chandra Research Institute, Chhatnag Road, Jhansi, Allahabad 211 019, India.
(Dated: February 2, 2008)

We study interacting scalar field theory non-minimally coupled to gravity in the FRW background. We show that for a specific choice of interaction terms, the energy-momentum tensor of the scalar field ϕ vanishes, and as a result the scalar field does not gravitate. The naive space dependent solution to equations of motion gives rise to singular field profile. We carefully analyze the energy-momentum tensor for such a solution and show that the singularity of the solution gives a subtle contribution to the energy-momentum tensor. The space dependent solution therefore is not non-gravitating. Our conclusion is applicable to other space-time dependent non-gravitating solutions as well. We study hybrid inflation scenario in this model when purely time dependent non-gravitating field is coupled to another scalar field χ .

PACS numbers: 04.20.Jb, 04.62.+v, 98.80.Cq

I. INTRODUCTION

Non-minimal coupling of the dilaton and the metric is a generic feature of any string theory (see e.g., [1]). While usual compactifications do not generate any potential, in the low energy limit, for the dilaton or other massless scalars coming from the string theory, it is generally believed that a potential is generated for them when one considers flux compactifications [2, 3]. Generation of potential for scalar fields in the flux compactification, makes them an arena for study of physics of the early universe. However, if one looks at models of inflation in these compactifications, a generic problem crops up in the slow roll inflation. One of the slow roll parameters, η , turns out to be sensitive to the size of compactified space and one finds that the slow roll condition, i.e., $\eta \ll 1$ is harder to achieve [4, 5, 6]. Smallness of the slow roll parameters ϵ and η ensures that the scale factor $a(t)$ in the Friedmann-Robertson-Walker (FRW) background grows exponentially in time or alternatively the Hubble parameter $H = a(t)^{-1} da(t)/dt$ is constant during the inflationary epoch [7].

The slow roll conditions come from the Einstein equations, where we demand that the energy-momentum tensor on the right hand side (RHS) of the equation is proportional to a constant times the FRW metric [7, 8, 9]. Since RHS contains the scalar field energy-momentum tensor this condition reflects upon the nature of the potential. However, if we have a non-minimally coupled scalar field with a potential such that it does not gravitate, i.e., its energy-momentum tensor vanishes identically then smallness of slow roll parameters ϵ and η becomes irrelevant. This is because throughout the rolling of the non-gravitating scalar field there is no contribution of it on the RHS of the Einstein equations. The shape

of the potential and rolling of the scalar field becomes relevant only when, by some other mechanism, it starts gravitating. If we achieve sufficient amount of inflation before the scalar field starts gravitating then subsequent time dependence of the scalar field has no bearing on the model of inflation. As we will see below, for a specific kind of non-minimal coupling, one gets a fixed potential for the scalar field so that it does not gravitate. It is not obvious if such a non-minimal coupling and/or the potential can be obtained from a generic flux compactification. Nevertheless inflation in the theory with a non-gravitating scalar field is a novel way of by-passing the ‘ η -problem’ [4] in the models of inflation in the flux compactifications [3].

If we are looking for a non-gravitating scalar field background, it could generically depend on the spatial coordinates as well. This question was addressed by [10] in the Minkowski spacetime and in the next section we will obtain similar results for FRW and anti-de Sitter backgrounds. As in [10, 11], we find that the classical non-gravitating solution allows space dependent scalar field profile in all these cases, including the Minkowski spacetime. However, in every case, this solution is singular. That is, the scalar field profile diverges at some point in space/spacetime. This singularity begs for a careful evaluation of the energy-momentum tensor in its neighbourhood. We analyze the energy-momentum tensor by regularizing the singular scalar field profile and find that there is a subtle contribution to the energy-momentum tensor from the singularity. This contribution is proportional to the Dirac δ -function but with an infinite multiplicative coefficient. Due to this contribution, the space dependent scalar field profile actually gravitates. While this rules out non-constant solution in the Minkowski spacetime, purely time dependent solution is allowed class of non-gravitating solutions in the FRW background. Similarly, purely radial dependent solution can be non-gravitating in the AdS background.

Since purely time dependent solution in the FRW background can be non-gravitating, we build a hybrid inflation model by coupling the non-gravitating solution to

^{*}E-mail: nabamita@mri.ernet.in

[†]E-mail: rajeev@mri.ernet.in

[‡]E-mail: dileep@mri.ernet.in

another scalar field χ , which is gravitating. Inflation is obtained by rolling of the non-gravitating scalar field and is exited by rolling of the gravitating scalar field, *i.e.*, χ . The rolling of χ is triggered by that of the non-gravitating field. Interesting feature of this model is that the non-gravitating scalar field starts gravitating as soon as χ starts rolling.

This paper is organized as follows. In section II, we review the result for non-gravitating scalar field and its interaction potential functional in the Minkowski background and obtain the same in FRW and AdS backgrounds. In section III, we perform the regularized energy-momentum tensor calculation in the vicinity of the singularity of the scalar field profile. We also mention the subtleties related to the issue of finding the backreacted metric in the presence of the non-vanishing stress tensor. In section IV, we propose a novel model of hybrid inflation by considering another scalar field χ . Finally, section V summarizes our results.

II. NON-GRAVITATING SCALAR FIELD IN VARIOUS BACKGROUNDS

In this section we will study a scalar field $\phi(\mathbf{x}, t)$ non-minimally coupled to gravity. We will first recall how the non-gravitating solution is obtained in the Minkowski space[10]. We will then obtain the solution in the FRW background as well as in the anti-de Sitter space. We are looking for a solution to the equations of motion such that the energy-momentum tensor for the field $\phi(\mathbf{x}, t)$ evaluated on this background vanishes. It imposes constraints which are sufficient to determine the form of the classical solution as well as the potential $V(\phi)$. The equation of motion of $\phi(\mathbf{x}, t)$ then becomes a consistency condition for the solution and the potential energy functional.

Let us consider a self-interacting scalar field $\phi(\mathbf{x}, t)$ non-minimally coupled to gravity with cosmological constant Λ in $(3+1)$ dimensions.

$$I_\phi = \int d^4x \sqrt{-g} \left(\frac{1}{2\kappa} (R + 2\Lambda) + \frac{1}{2} \phi \square \phi - \frac{1}{2} \xi R \phi^2 - V(\phi) \right). \quad (1)$$

The parameter ξ is a measure of non-minimality. In 3+1 dimensions, $\xi = 1/6$ leads to conformal coupling of the scalar field to the gravity and $\xi = 0$ corresponds to the usual minimal coupling. The field equation is

$$(\square - \xi R)\phi = V'(\phi), \quad (2)$$

where the additional term proportional to the Ricci scalar is a consequence of the non-minimal coupling. The conserved energy-momentum tensor is also modified due to the non-minimal coupling and is given by

$$\Theta_{\mu\nu} \equiv T_{\mu\nu} + \xi(g_{\mu\nu}\square - \nabla_\mu \nabla_\nu + \Lambda g_{\mu\nu})\phi^2, \quad (3)$$

where, $T_{\mu\nu}$ is the standard energy-momentum tensor for a minimally coupled scalar field,

$$T_{\mu\nu} = \partial_\mu \phi \partial_\nu \phi - g_{\mu\nu} \left(\frac{1}{2} \partial_\alpha \phi \partial^\alpha \phi + V(\phi) \right). \quad (4)$$

The expression for the energy-momentum tensor in eq.(3) is obtained in the following way. We first assume that in its original form, every component of the energy-momentum tensor $\Theta_{\mu\nu}$ vanishes. In that case the Einstein equation becomes $G_{\mu\nu} = \Lambda g_{\mu\nu}$. Using this we arrive at the form of $\Theta_{\mu\nu}$ given in eq.(3).

So far we have been looking at the general features of a scalar field non-minimally coupled to the gravity. Our aim is to look for a scalar field solution which does not gravitate in a given background. The solution is obtained by first doing a change of variable

$$\phi = \sigma^\alpha, \quad (5)$$

in the expression for the energy-momentum tensor (3) and writing them in terms of σ and its derivatives. Setting every component of $\Theta_{\mu\nu}$ to zero gives us a set of equations.

$$0 = \Theta_{\mu\nu} = \frac{(2\xi)^2}{(1-4\xi)} \frac{\phi^2}{\sigma} \nabla_\mu \partial_\nu \sigma, \quad \mu \neq \nu, \quad (6a)$$

$$0 = g_{ii}\Theta_{tt} - g_{tt}\Theta_{ii} = \frac{(2\xi)^2}{(1-4\xi)} \frac{\phi^2}{\sigma} (g_{ii}\nabla_t \partial_t \sigma - g_{tt}\nabla_i \partial_i \sigma) \text{ (no sum)}, \quad (6b)$$

$$0 = \Theta_{tt} = V(\phi) - \frac{(2\xi)^2 \phi^2}{(1-4\xi)\sigma} \left[\frac{\partial_\rho \sigma \partial^\rho \sigma}{2(1-4\xi)\sigma} - g^{ii} \sum_{i=1}^3 \nabla_i \partial_i \sigma \right] - \xi \phi^2 \Lambda. \quad (6c)$$

These equations impose constraints on σ . These equations are obtained by choosing $\alpha = 2\xi/(4\xi - 1)$, where $\xi \neq 0, 1/4$. While the case $\xi = 0$ is excluded because it takes us back to the minimally coupled scalar field theory, $\xi = 1/4$, can be treated separately and it can be shown that non-gravitating solution exists for $\xi = 1/4$ as well[10]. We will solve these constraints in various backgrounds below.

The Minkowski Space: Let us start with the Minkowski space background. The equations (6a) and (6b) are solved in the Minkowski space by[10]

$$\sigma(x^\mu) = a_1 x^\mu x_\mu + p^\mu x_\mu + a_2, \quad (7)$$

and the self interacting potential is given by

$$V(\phi) = \frac{2\xi^2}{(1-4\xi)^2} (\lambda \phi^{(1-2\xi)/\xi} + 48(\xi - \frac{1}{6}) a_1 \phi^{1/2\xi}), \quad (8)$$

where, $\lambda = p^\mu p_\mu - 4a_1 a_2$. We will get back to this solution in the next section.

The FRW Space: Let us now turn our attention to the FRW space. Background metric in this case is given by

$$ds^2 = -dt^2 + a^2(t)(dx^2 + dy^2 + dz^2) \quad (9)$$

where $a(t)$ is the scale factor of the spatial section of the background. By using the change of variables (5) and rearranging components of the energy-momentum tensor (3) appropriately we get three independent set of expressions, all of which are set to zero.

Eq. (6a) imposes constraints on the profile of σ . When neither μ nor ν is equal to t then the constraints on σ are such that we choose

$$\sigma = f(t)(X_1(x) + X_2(y) + X_3(z)). \quad (10)$$

If instead either μ or ν is equal to t then the constraint coming from the equation (6a) implies the function $f(t)$ is proportional to the scale factor $a(t)$, i.e.,

$$f(t) = c_0 a(t). \quad (11)$$

Linear combination of the diagonal components of the energy-momentum tensor in the eq.(6b) determines the solution completely. That is, while on one hand, it can be used to obtain behaviour of the scale factor, on the other hand, it also determines the functional forms of $X_1(x)$, $X_2(y)$ and $X_3(z)$ up to constants of integration. This equation also requires exponential behaviour of the scale factor. This is consistent with the Einstein equation which can be written in terms of the Hubble parameter $H = a(t)^{-1}(da(t)/dt)$ and the cosmological constant Λ as,

$$3H^2 = -\Lambda, \quad a(t) = a_0 \exp(Ht). \quad (12)$$

Vanishing of the double spatial derivative of σ , in eq.(6b), implies that the functions X_i are at most linear in their arguments for all i , for example,

$$X_1(x) = p_1 x + c_1. \quad (13)$$

Using eqs.(10-13) in the relation (5) we get

$$\phi(\mathbf{x}, t) = \sigma^{2\xi/(4\xi-1)} = (c_0 a(t)(p_i x^i + c))^{2\xi/(4\xi-1)}, \quad (14)$$

where, $c = c_1 + c_2 + c_3$ and $p_i x^i = p_1 x + p_2 y + p_3 z$. We can now determine the potential $V(\phi)$ using the Θ_{tt} equation and the solution (14).

$$V(\phi) = \left[\frac{2\xi^2}{(4\xi-1)^2} (5 - 24\xi) - 3\xi \right] H^2 \phi^2 + \frac{2\xi^2}{(4\xi-1)^2} \left(c_0^2 \sum_{i=1}^3 p_i^2 \right) \phi^{(1-2\xi)/\xi}. \quad (15)$$

Let us first notice that for all $\xi < 1/4$, second term in the potential is dominant for large ϕ and if ξ is such that the potential $V(\phi)$ is even under $\phi \rightarrow -\phi$ then the potential is bounded from below. Since the first term is negative, for the parity symmetric cases we have a double well potential(fig.1(a)). The solution (14) obtained by solving the energy-momentum tensor constraints also satisfies the ϕ equation of motion. This consistency condition ensures that the energy-momentum tensor is conserved in

this background. This solution diverges for large x^i when $\xi > 1/4$ ($\alpha > 0$), on the other hand, for $\xi < 1/4$ ($\alpha < 0$) it diverges at a point where $(p_i x^i + c) = 0$ (fig.1(b)). We choose ξ in such a way that α , which in general is a fraction, has odd denominator. This ensures that at least one real solution always exists. The classical solution $\phi(\mathbf{x}, t)$ depends on the scale factor $a(t)$ and when $\xi < 1/4$, the field $\phi(\mathbf{x}, t)$ is inversely proportional to a ξ dependent power of $a(t)$. Thus, in addition to having the space dependent profile, the amplitude of the classical solution depends on the scale factor and decreases rapidly for $\xi < 1/4$.

For the sake of completeness let us consider the case of $\xi = 1/4$. In this case we carry out analysis similar to that of [10] and obtain the classical solution $\phi(\mathbf{x}, t)$ and the potential $V(\phi)$ exactly.

$$\phi(\mathbf{x}, t) = \exp(c_0 e^{Ht} (p_i x^i + c)), \quad (16)$$

$$V(\phi) = \phi^2 \left[\sum_{i=1}^3 \frac{(c_0 p_i)^2}{2} - \frac{3H^2}{4} - \frac{(H \ln \phi)^2}{2} - \frac{3H^2}{2} \ln \phi \right]. \quad (17)$$

The Anti-de Sitter space: Let us now turn our attention to the Anti-de Sitter space(AdS). The metric on the four dimensional AdS space is given as,

$$ds^2 = -\frac{r^2}{b^2} dt^2 + \frac{b^2}{r^2} dr^2 + \frac{r^2}{b^2} (dx^2 + dy^2), \quad (18)$$

where, b is the radius of the anti-de Sitter space. The conditions coming from the $\Theta_{\mu\nu}$ ($\mu \neq \nu$) can be satisfied by choosing the following ansatz for σ

$$\sigma = a(r)[f_1(t) + f_2(x) + f_3(y)]. \quad (19)$$

Diagonal components of the energy-momentum tensor then give

$$a(r) = Cr, \quad (20)$$

where C is the integration constant and

$$\sigma = Cr(t + x + y + d), \quad (21)$$

where d is another constant of integration. The potential is obtained by solving $\Theta_{tt} = 0$.

$$V(\phi) = \alpha^2 b^2 C^2 \phi^{\frac{(1-2\xi)}{\xi}} + (8\xi\alpha + 5\xi + \frac{\alpha^2}{2}) \frac{\phi^2}{b^2}. \quad (22)$$

It is worth emphasizing that this solution solves the equation of motion ensuring conservation of the energy-momentum tensor.

While we have managed to obtain non-gravitating solutions to the equations of motion in various backgrounds, all these solutions are singular. If one chooses $\xi > 1/4$ then the solution diverges at infinity and if $\xi < 1/4$ then




FIG. 1: Fig.(a) shows generic form of the potential for the non-gravitating field ϕ in the FRW background. Curve in fig.(b) denotes ϕ^2 , square of the classical solution at a fixed time.

it diverges for some finite value of space-time coordinates. This divergence in the solution has a subtle contribution to the energy-momentum tensor. In the next section we will carefully analyze the solution and show that naive conclusion of vanishing of the energy-momentum tensor is flawed.

III. REGULARIZED ENERGY-MOMENTUM TENSOR

In this section we will show that the naive computation of the stress tensor for the “non-gravitating” solution in the previous section fails in the vicinity of the singularity. Singular nature of the solution, depending on the choice of ξ , leads to either discontinuous profile for ϕ or its derivative $\partial_\mu \phi$. Since the energy-momentum tensor contains terms upto second derivative of ϕ , this discontinuity can lead to a potential δ -function contribution. To analyze this we will regularize the solution in the neighbourhood of singularity. We will illustrate this method for the Minkowski space as well as for the FRW space, a similar analysis carries through for the anti-de Sitter space. For concreteness, we will carry out our analysis for $\xi < 1/4$, but it can be easily extended to $\xi > 1/4$. While in the former case the singularity is in the finite domain, it is at infinity for the latter case.

The Minkowski Space: In this case we regularize solution by regularizing the function σ as follows

$$\sigma = \sqrt{(x+t)^2 + \epsilon^2} \sqrt{(x-t)^2 + \epsilon^2}, \quad (23)$$

where, for illustration, we have chosen $a_1 = 1$, $p_\mu = 0$ and $a_2 = 0$ in (7). It is evident that as $\epsilon \rightarrow 0$, we recover original form for σ , namely $\sigma = x^2 - t^2$. Notice that we are choosing the profile for σ such that it depends only on one spatial dimension. This suffices to illustrate our point and reinstating its full space dependence does not alter our conclusion. Using this regularized form of σ , and therefore ϕ , we can write down any component of the energy-momentum tensor $\Theta_{\mu\nu}$ at an arbitrary point in Minkowski space. The non-vanishing off-diagonal com-

ponent of $\Theta_{\mu\nu}$ is,

$$\Theta_{tx} = -2\xi\alpha\sigma^{2\alpha} \left[\frac{\epsilon^2}{((x+t)^2 + \epsilon^2)^2} - \frac{\epsilon^2}{((x-t)^2 + \epsilon^2)^2} \right]. \quad (24)$$

At any point where the field profile ϕ is finite, we can show using independent methods that every component of the energy-momentum tensor vanishes. We, therefore, need to restrict ourselves to the neighbourhood of the singularity. Taking $(x+t) \sim \epsilon$ and $(x-t) \sim \epsilon$,

$$\Theta_{tx} = -2\xi\alpha\epsilon^{4\alpha} f_1 \left(\frac{x}{\epsilon}, \frac{t}{\epsilon}, \epsilon \right). \quad (25)$$

As expected the function f_1 is such that when $x, t \gg \epsilon$ it vanishes as $\epsilon \rightarrow 0$, and when $x, t \sim \epsilon$, $f_1 \sim \epsilon^{-2}$. Therefore, in the neighbourhood of the singular point, i.e., when $x, t \sim \epsilon$

$$\Theta_{tx} \sim \epsilon^{(4\alpha-2)}. \quad (26)$$

Since $4\alpha - 2$ is negative for $\xi < 1/4$, we see that the Θ_{tx} component of the energy-momentum tensor is divergent in the limit $\epsilon \rightarrow 0$. In fact, analysis of diagonal components of the energy-momentum tensor shows that they also diverge as $\epsilon^{(4\alpha-2)}$ in the neighbourhood of the singular point. Notice that in the limit $\epsilon \rightarrow 0$ the function f_1 has δ -function support at the location of the singularity, however, the coefficient multiplying this δ -function diverges as we remove the regulator. This severe singular contribution to $\Theta_{\mu\nu}$ invalidates the naive derivation of the non-gravitating solution in the previous section.

The FRW Space: The situation is not any better in the FRW background. We write the regularized solution as

$$\phi = a(t)^\alpha x(x^2 + \epsilon^2)^\beta. \quad (27)$$

where $\beta = (\alpha - 1)/2$. Substituting this regularized form of the solution in the energy-momentum tensor, we find that all off diagonal components of $\Theta_{\mu\nu}$ vanish identically at the singular point. The diagonal components, on the other hand, take the form

$$\Theta_{tt} = a(t)^{2\alpha} \epsilon^{4\beta} f_2 \left(\frac{x}{\epsilon} \right). \quad (28)$$

The function f_2 vanishes when $x \gg \epsilon$ and $\epsilon \rightarrow 0$ and when $x \sim \epsilon$, it is constant. Therefore near the singular point *i.e.* when $x \sim \epsilon$ and $\epsilon \rightarrow 0$, Θ_{tt} diverges as $\epsilon^{4\beta}$. All other diagonal components exhibit the similar divergent behaviour,

$$\Theta_{ii} = a(t)^{2\alpha+2} \epsilon^{4\beta-2} f_2\left(\frac{x}{\epsilon}\right). \quad (29)$$

This divergence is as severe as in the case of the Minkowski space. Analysis of the energy-momentum tensor of the scalar field in the anti-de Sitter space proceeds in the similar fashion as in the FRW case.

We can now ask what is the back reaction of this contribution to the energy-momentum tensor on the background metric. Notice that in the FRW case the Dirac δ -function contribution exists only for the diagonal components of the energy-momentum tensor and all the off-diagonal components continue to vanish. With this in mind we consider the metric ansatz for the backreacted geometry to be

$$g_{\mu\nu} = \text{diag}(-f_1(x, t), f_2(x, t), f_3(x, t), f_3(x, t)). \quad (30)$$

Substituting this ansatz in the Einstein equation gives us a set of coupled non-linear inhomogeneous partial differential equations. To make things even harder, we find, coefficient of the inhomogeneous term is time dependent. In order to seek a solution we will attempt a separable ansatz for the solution. We will also look at the solution on one side of the singularity. A similar solution on the other side of the singularity can then be matched with the first one along with the contribution from the singularity. Since the energy-momentum tensor vanishes away from the singularity, RHS of the Einstein equation is proportional to the cosmological constant term only. As an illustration let us consider an ansatz

$$f_i(x, t) = (cx + d)^{\alpha_i} g_i(t). \quad (31)$$

With this ansatz, we get one sided solution with $\alpha_1 = \alpha_3 = 0$ and $\alpha_2 = -2$ and two classes of time dependent terms are $g_1(t) = c_1$, $g_2(t) = c_2 e^{-2t}$, $g_3(t) = c_3 e^t$ or $g_1(t) = c_1$, $g_2(t) = c_2 e^{2t}$, $g_3(t) = c_3 e^{-t}$ where c_1 , c_2 and c_3 are integration constants. We can now take the δ -function contribution into account to match solutions on either side of the singularity. Due to the time dependent coefficient of the δ -function, we end up with inconsistent powers of the scale factor on the right and left hand side of the Einstein equation. We therefore conclude that the separable ansatz does not give a solution to our coupled inhomogeneous PDEs. We have tried out more general separable ansatzes but all of them have suffered the same fate.

Since vanishing of the energy-momentum tensor also gives rise to coupled nonlinear, but homogeneous, partial differential equations, we can attempt to get another solution of these equations which are not singular. This does not seem to lead to any new solution. Noticing the fact that the scalar field profile is proportional to

$\exp(Ht)$, we can try a field redefinition, $\phi = \exp(\psi)$ and attempt to seek a solution

$$\psi(x, t) = At + f(x). \quad (32)$$

This ansatz gives us back the singular solution. Instead of the additive ansatz if we take a multiplicative ansatz, *i.e.*, $\psi(x, t) = At f(x)$ we end up with a contradiction. While one equation demands the scale factor to be a function of time only, other equation is satisfied only if the scale factor has spatial dependence. Replacing At in the product ansatz by any other time dependent function does not alter the result.

IV. HYBRID INFLATION MODEL

In the previous section, we saw that space dependent scalar field profile fails to be non-gravitating. However, if we look at purely time dependent scalar field profile in the FRW background, it does not lead to any subtle contribution to the energy-momentum tensor. We can therefore use the space independent scalar field profile to construct a model of inflation.

To do that we will couple another scalar field $\chi(\mathbf{x}, t)$ to the model studied above. For simplicity we consider minimal coupling between $\chi(\mathbf{x}, t)$ and gravity. The additional term in the action corresponding to the field $\chi(\mathbf{x}, t)$ is

$$I_\chi = \int d^4x \sqrt{-g} \left(\frac{1}{2} \chi \square \chi - V_\chi(\phi, \chi) \right). \quad (33)$$

We choose the interaction term V_χ as

$$V_\chi(\phi, \chi) = \frac{\lambda_1}{2} (\phi^2 - \phi_0^2) \chi^2 + \lambda_2 \chi^4, \quad (34)$$

where, λ_i are positive for $i = 1, 2$. We define the total action to be $I = I_\phi + I_\chi$. We can now justify the cosmological constant term in I_ϕ by interpreting it as the value of the potential V_χ at $\chi = 0$. This essentially amounts to redefining value of V_χ at $\chi = 0$. For the model to be studied here, we will choose $\xi < 1/4$ therefore, in this case magnitude of the classical solution $\phi(\mathbf{x}, t)$ decreases as a function of time as the scale factor $a(t)$ grows.

For a fixed ϕ_0 , it is easy to see from the potential (34), that we have two regimes corresponding to $\phi > \phi_0$ and $\phi < \phi_0$. As we discussed in the previous section, space dependent profile for ϕ is singular and it gives rise to a subtle contribution to the energy-momentum tensor. The space dependent solution (14) therefore is not a bona fide non-gravitating solution. Purely time dependent configuration for ϕ , however, does not suffer from this problem and is a legitimate non-gravitating field configuration. For $\phi > \phi_0$, the potential for the field χ has a unique minimum at $\chi = 0$ and in this case the field ϕ does not gravitate. The exponential growth of scale factor is then determined solely by the cosmological constant $\Lambda = \langle V_\chi \rangle|_{\chi=0}$ and the rate at which the field ϕ is rolling

is irrelevant. In particular, the slow roll parameters ϵ and η determined from the potential $V(\phi)$ have no bearing on the nature of the inflation as $\phi(\mathbf{x}, t)$ is not gravitating in this domain. This situation continues until $\phi(\mathbf{x}, t_0) = \phi_0$ at some time $t = t_0$. For all subsequent times after t_0 , we are in a situation where $\phi < \phi_0$. A sufficient amount of inflation can be obtained by adjusting initial value of ϕ .

Once $\phi < \phi_0$, the field χ develops tachyonic instability at $\chi = 0$. This instability makes χ roll towards the true minimum. The rolling of χ affects exponential growth of the scale factor. This is because two new terms are generated due to rolling of the field χ . First of all, rolling of χ brings in I_χ into the picture, *i.e.*, the field χ contributes to the energy-momentum tensor and consequently to the right hand side of the Einstein equation. Secondly, non-zero vacuum expectation value of χ , changes the potential $V(\phi)$ by adding a ϕ^2 term which modifies mass of ϕ . Due to this additional term the field ϕ ceases to be a non-gravitating field and starts contributing to the right hand side of the Einstein equation. The first term is identical to the scalar field in the hybrid inflation models, which is responsible for exit from the inflationary regime. It would be interesting to study this model further and check its validity against currently available CMB data[12, 13].

A. Conformal Transformation

It is known for some time that action of the scalar field nonminimally coupled to gravity can be brought into canonical minimally coupled form by conformal transformation(see *e.g.*, [14]). Consider the action

$$S = \int d^4x \sqrt{-g} \left[\frac{\Omega^2}{2\kappa} R - \frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - V(\phi) \right], \quad (35)$$

where, $\Omega^2 = 1 - \kappa\xi\phi^2$. A minimally coupled scalar field theory can be obtained by doing the conformal transformation,

$$\Omega^2 g_{\mu\nu} = \tilde{g}_{\mu\nu}, \quad (36)$$

and followed by the redefinition of the scalar field

$$d\tilde{\phi} = \frac{[1 - \kappa\xi(1 - 6\xi)\phi^2]^{1/2}}{\Omega^2} d\phi. \quad (37)$$

The new action written in terms of $\tilde{\phi}$ with the potential energy functional

$$\tilde{V}(\tilde{\phi}) = \frac{V(\phi)}{\Omega^4}, \quad (38)$$

is minimally coupled to gravity. The equation (37) can be integrated to write $\tilde{\phi}$ as a function of ϕ

$$\tilde{\phi} = \frac{1}{\sqrt{2\kappa\xi}} \left(2\sqrt{3}\xi \tanh^{-1} \left(\frac{\sqrt{6\kappa\xi}\phi}{\sqrt{1 - \kappa\xi(1 - 6\xi)\phi^2}} \right) - \sqrt{2\xi(-1 + 6\xi)} \sinh^{-1}(\sqrt{\kappa\xi(-1 + 6\xi)}\phi) \right). \quad (39)$$

Substituting this redefinition in the action gives rise to a pretty complicated potential energy functional. Nevertheless, in principle, it is possible to get non-gravitating scalar field background even in the minimally coupled case. This result holds for all backgrounds including the Minkowski space[10]. The non-minimally coupled situation is, clearly, much easier to deal with.

In case of the hybrid inflation model, we have an added complication in the minimally coupled situation. This is due to the fact that the conformal transformation which relates the non-minimally coupled theory to the minimally coupled one shows up in front of the kinetic energy term of χ as well as the potential energy term. The action for χ becomes

$$\mathcal{L}_\chi = -\Omega^{-2} \tilde{g}^{\mu\nu} \partial_\mu \chi \partial_\nu \chi - \Omega^{-4} V_\chi(\tilde{\phi}, \chi). \quad (40)$$

Since Ω is field dependent, it generates a non-trivial metric in the field space.

V. DISCUSSION AND SUMMARY

We have considered a novel model for inflation with a non-minimally coupled non-gravitating scalar field. Naive non-gravitating classical solution for the scalar field ϕ has space dependent profile. Such a space dependent profile is singular. This singularity gives rise to a subtle but divergent contribution to the energy-momentum tensor. Due to this contribution, the space dependent profile of ϕ fails to be a non-gravitating solution. The divergent contribution to the energy-momentum tensor has a delta function support at the location of singularity of the classical solution. The divergence is more severe than just the delta function because for any $\alpha \neq 0$ the coefficient multiplying the delta function also diverges. This problem is not specific only to the space dependent profile of the scalar field in the FRW background. This problem exists for solutions in the Minkowski space and the anti-de Sitter space as well. We attempted to determine backreaction of this divergent contribution on the geometry. This gives rise to a set of coupled inhomogeneous non-linear partial differential equations. Several obvious and not-so-obvious ansatz fail to solve these equations.

Space independent profile for ϕ , however, does retain the non-gravitating property. We use this fact to write down a hybrid inflation model by coupling ϕ to another scalar field χ . Utility of this model is in the non-gravitating nature of the scalar field ϕ . Due to this property the inflation is completely driven by the cosmological constant, which is the value of the potential $V_\chi(\phi, \chi)$ at $\chi = 0$. Expansion of the universe is independent of how ϕ rolls down and enters into the gravitating regime. Value of slow roll parameters ϵ and η is therefore of little consequence. Although non-minimally coupled scalar field can, after a conformal transformation, be brought into minimally coupled form, we show that

such a field redefinition is not very convenient. It is, in fact, more efficient to work with non-minimal coupling.

We would like to thank J. S. Bagla, S. Dutta, R. Gopakumar, T. Padmanabhan and L. Sriramkumar for numerous discussions. We would especially like to thank

A. Sen for many an interesting discussions and for emphasising on the regularized energy-momentum tensor near the singularity. We would also like to thank D. Choudhury for useful correspondence and pointing out subtleties involved in seeking numerical solutions to coupled inhomogeneous PDEs.

-
- [1] J. Polchinski, “String theory. Vol. 1: An introduction to the bosonic string,” *Cambridge, UK: Univ. Pr. (1998) 402 p.*
 - [2] J. Polchinski, “String theory. Vol. 2: Superstring theory and beyond,” *Cambridge, UK: Univ. Pr. (1998) 531 p.*
 - [3] M. R. Douglas and S. Kachru, “Flux compactification,” arXiv:hep-th/0610102.
 - [4] J. McDonald, “F-term hybrid inflation, the eta-problem and extra dimensions,” *JHEP* **0212**, 029 (2002) [arXiv:hep-ph/0201016].
 - [5] A. Riotto, “Inflation and the theory of cosmological perturbations,” arXiv:hep-ph/0210162.
 - [6] R. Holman and J. A. Hutasoit, “Axionic inflation from large volume flux compactifications,” arXiv:hep-th/0603246.
 - [7] A. Linde, “Particle Physics and Inflationary Cosmology,” *Harwood Academic, Chur, Switzerland, (1990) 362 p.*, arXiv:hep-th/0503203; A. R. Liddle and D. H. Lyth, “Cosmological Inflation and Large-Scale Structure,” *Cambridge, UK: Univ. Pr. (1999) 400 p.*
 - [8] B. A. Bassett, S. Tsujikawa and D. Wands, “Inflation dynamics and reheating,” *Rev. Mod. Phys.* **78**, 537 (2006) [arXiv:astro-ph/0507632].
 - [9] V. Mukhanov, “Physical foundations of cosmology,” *Cambridge, UK: Univ. Pr. (2005) 421 p.*
 - [10] E. Ayon-Beato, C. Martinez, R. Troncoso and J. Zanelli, “Nonminimally coupled scalar fields may not curve spacetime,” *Phys. Rev. D* **71**, 104037 (2005)
 - [11] D. A. Demir, B. Pulice, “Non-Gravitating Scalars and Spacetime Compactification,” *Phys. Lett. B* **638**, 1 (2006).
 - [12] D. N. Spergel *et al.*, “Wilkinson Microwave Anisotropy Probe (WMAP) three year results: Implications for cosmology,” [arXiv:astro-ph/0603449].
 - [13] R. Jeannerot and M. Postma, “Confronting hybrid inflation in supergravity with CMB data,” *JHEP* **0505**, 071 (2005) [arXiv:hep-ph/0503146].
 - [14] V. Faraoni, E. Gunzig and P. Nardone, “Conformal transformations in classical gravitational theories and in cosmology,” *Fund. Cosmic Phys.* **20**, 121 (1999) [arXiv:gr-qc/9811047].